

10/540086
JC17 Rec'd PCT/PTO 20 JUN 2005

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims

Claims 1-24 (cancelled)

Claim 25 (new): Oligonucleotide primers for specific amplification of a *hupB* gene of *Mycobacterium* species selected from the group consisting of Seq ID No. 1, Seq ID No. 2, Seq ID No. 3, Seq ID No. 4 and Seq ID No. 5.

Claim 26 (new): A method for differentiating *Mycobacterium* species based on target gene encoding for histone like proteins such as *hupB* comprising:

- a) obtaining DNA from culture or from clinical samples;
- b) amplifying a part of the target gene encoding for histone like proteins such as *hupB* of *Mycobacterium* species using said DNA as a template in a polymerase chain reaction with a pair of oligonucleotide primer according to claim 25; and
- c) detecting said amplified fragment of the *hupB* gene to detect the presence of *Mycobacterial* species or not and differentiating *Mycobacterium tuberculosis* from *Mycobacterium bovis* based on the size of the amplified fragment.

Claim 27 (new): The method according to claim 26, wherein said *Mycobacterium tuberculosis*, and or *Mycobacterium bovis* species is selected from a group of genetically related *Mycobacteria* and from unrelated microorganisms.

Claim 28 (new): The method according to claim 26, wherein the pair of oligonucleotide primers comprises Seq ID No. 1 and Seq ID No. 2.

Claim 29 (new): The method according to claim 26, wherein the pair of oligonucleotide primers comprises Seq ID No. 3 and Seq ID No. 2.

Claim 30 (new): The method according to claim 26, wherein the pair of oligonucleotide primers comprises Seq ID No. 4 and Seq ID No. 5.

Claim 31 (new): The method according to claim 26, wherein the amplified fragments are detected by ethidium bromide staining or DNA probe hybridization.

Claim 32 (new): The method as claimed in claim 26, wherein the step of differentiating comprises the steps of:

- a) designing a set of primers selected from the group consisting of Seq ID No. 1, Seq ID No. 2, Seq ID No. 3, Seq ID No. 4 and Seq ID No. 5 to amplify a part of said *hupB* gene from *Mycobacterium tuberculosis* and *Mycobacterium bovis*;
- b) obtaining DNA from culture or from clinical samples;
- c) amplifying a part of the target gene encoding for histone like proteins such as *hupB* of *Mycobacterium* species using said DNA as a template in a polymerase chain reaction with a pair of said oligonucleotide primers;
- d) analyzing and validating the size of the amplified fragment;
- e) determining the complete sequence of said amplified fragments; and
- f) inferring from the sequence whether it is *Mycobacterium tuberculosis* or *Mycobacterium bovis*.

Claim 33 (new): The method according to claim 31, wherein the DNA probe consists of Seq ID No. 7 or Seq ID No. 8 or a complement thereof tagged with a detectable label.

Claim 34 (new): The method as claimed in claim 26, wherein the step of differentiation consists in determining the smaller size of the amplified fragment obtained from *Mycobacterium bovis*.

Claim 35 (new): The method according to claim 28, wherein the PCR amplified fragment in *Mycobacterium bovis* is 618 bp.

Claim 36 (new): The method according to claim 28, wherein the PCR amplified fragment in *Mycobacterium tuberculosis* is 645 bp.

Claim 37 (new): The method according to claim 29, wherein the PCR amplified fragment in *Mycobacterium bovis* is 291 bp.

Claim 38 (new): The method according to claim 30, wherein the PCR amplified fragment in *Mycobacterium tuberculosis* is 318 bp.

Claim 39 (new): The method according to claim 29, wherein the PCR amplified fragment in *Mycobacterium bovis* is 89 bp.

Claim 40 (new): The method according to claim 29, wherein the PCR amplified fragment in *Mycobacterium tuberculosis* is 116 bp.

Claim 41 (new): The method according to claim 26, wherein the PCR amplified fragment in *Mycobacterium bovis* is 27 bp smaller than that of *Mycobacterium tuberculosis*.

Claim 42 (new): The method as claimed in claim 26, wherein differentiating *Mycobacterium tuberculosis* and *Mycobacterium bovis* comprises the steps of:

- a) amplifying a part of the target *hupB* gene from *Mycobacterium tuberculosis* and *Mycobacterium bovis* in a polymerase chain reaction with primers Seq ID No. 1 and Seq ID No. 2;
- b) restricting the amplified fragment with *Hpa II* restriction enzyme to produce restricted fragments;
- c) separating the restricted fragments by electrophoresis on 12% polyacrylamide gel; and
- d) detecting the restricted fragments by staining with ethidium bromide.

Claim 43 (new): The method according to claim 42, wherein the restricted fragment in *Mycobacterium tuberculosis* is 280 bp and 150 bp.

Claim 44 (new): The method according to claim 42, wherein the restricted fragment in *Mycobacterium bovis* is 253 bp and 150 bp.

Claim 45 (new): A *hupB* gene comprised of Seq ID No. 8 as claimed in claim 25.

Claim 46 (new): A *hupB* gene comprised of Seq ID No. 7 as claimed in claim 25.

Application No. Not Yet Assigned
Paper Dated: June 20, 2005
In Reply to USPTO Correspondence of N/A
Attorney Docket No. 4544-051936

Customer No. 28289

AMENDMENTS TO THE ABSTRACT

After the claims, please insert a page containing the Abstract Of The Disclosure, which is attached hereto as a separately typed page.